UTAH FIELD ORNITHOLOGISTS SIGNIFICANT SIGHTING RECORD DOCUMENTATION

1982-22 afo

TO THE PART OF THE	
DO NOT WRITE IN THIS SPACE FOR U.F.O. RECORDS AND D	
Latilong Reporter	Sci Name
Habitat Record type	Serial No. JAN 5 1 PEND (1982-003).
Date/timeDisposition	Cx Nos.
Common Name Oldsquaw	Weather at time of this observation. 75 per cent overcast; 34°F; wind S at 5-7 mph; water open
Race/Sub-species Clangula hyemalis	
Sight date 01/30/82 time 1100 MST duration 10 mins.	
Distance from bird 100-300 maters (estimated)	Prior weather & number of days since chang A weak low pressure trough had passed the previous night. Last "major" storm 4 days earlier on 26/27th.
Light conditions good, overcast sky	
Optical equip. 48x50 Bushnell Sentry II scope	
Photo. equip. None	
Number of this species seen 2	Locality
	Salt Lake State Park, Davis County, Utah. About 9.3 miles WSW of Syracuse USPO.
Male ☐ Adult ☐k Age unk. Female ☐ Subadult ☐ + or Imm Sex Unk. ☐ Juv/Imm ☐ 2/3 year bird	☐ Breeding ☐ Dark Phase ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐
Description of bird.Size, Colors, all field marks O	BSERVED -Unique features of this bird, if any.

Two birds of stender buck shape were seen which reminded me somewhat of small pintails. The birds were near some common joidsnayes and appeared to be slighter and a little smaller. In general, both birds were quite light, exhibiting a great deal of white in their plumage. Ventrally both were white and the neck and head were generally white on both. The bills of both birds were generally gray but appeared to have an orange wash; profile bill shape was rounder than that of most ducks.

Plumaça details show one bird to be male and the other female. The alleged male bird had a white forehead, crown and mape, with a dark back and a sort of star-like white pattern on the folial wings. The facial markings included a gray, oval-shaped curicular patch with a small darker oval patch inside the larger patch at its rear. This bird did not have the longer tail feathers of tais plumage.

The alleged female bird had a dingy gray crown and mape and a much simpler gray auricular patch which appeared to blend in with the mape gray area. The bird was dingier appearing on the back than the male and did not have significant markings on its folded wings.

marks which were not seen. Why not?

terson and Robbins indicate three field marks nich I did not see:

the "needle-like" tail of the male. (1)

- (2) the contrast of all dark wings to white body (birds did not fly).
- (3) the dark breast of the male bird.

Variations from expected field marks.

No variation seen.

List similar species and describe why or how you eliminated them.

To my knowledge the only similiar species are pintail (dark head markings on white neck), and bufflehead, scaups, hooded merganser (extensive white on body). The white crown and the facial patterning eliminates pintail, scaup and goldeneyes while the white crown and flanks eliminates the merganser.

Behavior of this bird. Include names of other birds with this one and any interaction. Birds were swimming together in the harbor nearby but not associating with a small flock of common goldeneye. Birds swimming was steady and even. They swam towards the outlet of the harbor to the Great Salt Lake.

Habitat this location and near by or adjacent habitat if significant.

Protected salt water harbor adjacent to large bodies of both fresh and salt water in the Great Salt Lake. Landward, Antelope Island is sagebrush community.

Describe birds song or sounds it made during this obser-Birds were silent during observation.

Names of other observers.A/C+phon

What is your experience with this or similar birds?

I feel myself to be generally competen, with the ducks. I have seen basic (winter) plumage oldsquaw on several occasions in Utah and California.

Print name & mailing address bel

Joshua A. Burns 1572 South 1400 East Salt Lake City, Utah 84105

A/C and Phone No.801-484-7260