DO NOT WRITE IN THIS SPACE. FOR UTAH BIRD RECORDS CODING ONLY. Latilong Record type		Serial No		
Habitat		Disposition	2 (95)	
Common Name <u>Hooded Warbler</u> Scientific Name Wilsonia citrina Obs. date Nov. 5, '87me(s) 1:10-1:20 p.m.			Weather at time of this observation. Slightly overcast with periods sun; temps. in 60's or low 70'	
Duration of obs. about 10 min. Distance from bird $10-20$ yards Light conditions scattered clouds; muted light Optical equip. $7x50$ Tasco binoculars Photo equip. none Notes made at time of sighting x Date report prepared			Prior weather and number of days since change Warm weather for two-three weed least warm for this time of years no rain or snow; low temps. In Locality About 1½ miles south of Pintum in Washington County and about east of the I-15 freeway in pri	ear. high ca, Uta k mil lnion-
Notes made later from med Enter number of birds in each category. Total number observed.		AGES(S) Adults × Juv/Imms 2/3 yr birds	juniper and live oak community PLUMAGE(S) Breeding X Transition Avaliable Teclipse Dark phase Enclosed Winter Light ph. Other	none

Description of bird: Size, colors, all field marks <u>OBSERVED</u>. - Unique features of <u>this</u> individual, if any. Size- Small, warbler size; about 5-6 inches in lenght.

Colors

Abdomen and flanks brilliant yellow; face also, Yellow was similar to a Wilson's Warbler color.

Head was all black: black continued underneath in a continuous band across th upper breast until it merged with the black on the opposite side of head form a complete hood.

Back was olive-brown with very faint but noticeable brown lines or streaks. When the bird was in flight whitish outer tail feathers were observed; not as striking as in juncos, but very noticeable in contrast to darker brownish central rectrices.

Eyes were dark.

pecel our

Feet were light colored, but didn't see them well.

I should mention that I was hunting quail (I had observed a large covey of about 50-70 birds previously in that vicinity), so I was carrying a 20 gauge shotgun as well as my binoculars. The urge to collect the bird was tempered by the fact that had I shot it at the distance at which I was observing it (10-20 yards there would not have been much left to identify. I am satisfied that the identification is correct, I just do not have a specimen to prove it.

Field marks which were <u>not</u> seen. Why not?	Variations from expected field marks.
that Saw all the fieldguides list as important for positive identifi- cation.	none
List similar species and describe why or how you eliminated t	them.
Wilson's Warbler is the only one in te be confused with this species being descr and the black across the chest as well as as a possibility.	ibed. The amount of black on the head
Behavior of this bird. Other species seen with this one. An	y interaction between birds?
This bird was alone. Did not see it i	
	not on the outside or on top but into vation difficult. When it flew into the low in the branches and in the interious moving about in warbler fashion When it was between 5-10 feet high i
Habitat at this location. Adjacent or near by habitat, if si	gnificant.
Habitat was pinion-juniper with scatted I-15 freeway was between 1-15 mile to to the east. The substratum was lavatery little grass or forb.	ered thick clumps of shrub live oak. the west and Ash Creek was about 1/2 mile rock (scattered) and reddish dirt with
Describe bird's song or sounds.	Print reporter's name, address, phone number
The bird only gave short, sharp chip notes. And not many at that.	Merrill Webb 1063 East 400 North Orem, Utah 84057
	801-224-6113
	Signature Merrill Webb
What is your experience with this or similar birds? No prior experience with this species;	Corroborating observers not reporting separately.
a lot of experience with western species of warblers (about 20 years worth).	Print Name Signature
	Print Name Signature
	Print Name Signature
	Print Name Signature

Variations from expected field marks.