UTAH FIELD ORNITHOLOGISTS UTAH BIRD RECORDS COMMITTEE

DO NOT WRITE IN THIS SPACE	E. FOR UTAH BI	RD RECORDS CODING ONLY.	- I	*
Latilong		Record type	Serial No. <u>84</u> -	-001
Habitat	200 C C C C C C C C C C C C C C C C C C	Disposition	Cx Nos	
Common Name Glaucous Gull 3			Weather at time of this observation.	
Scientific Name Larus hyperboreus			cloudy, overcast, occasional drizzle	
Obs. date 1/24/84 Time(s)2:30-4:00 PM				
Juration of obs. 15 mins, then later 2 mins.			Prior weather and number of days since change very cold with valley fog	
distance from bird less than 200 ft.			very cold with valley rog	
Light conditions fair-poor, grayish day				1 N N
Optical equip. 9X binoculars			Locality	
Photo equip.			Salt Lake Garbage I	Dump
Notes made at time of sighting Date report prepared		A Compress CA		
Notes made later from memory x 1/28/84				
Enter number of birds in each category.	GENDER(S)	AGES(S)	PLUMAGE(S)	PHOTO(S)
	Males	Adults	Breeding Transition	Avaliable
Total number observed.	Females	Juv/Imms 1	Eclipse Dark phase	Enclosed
11	for Imms 1	2/3 yr birds	Winter 1-1stight ph	
	Unknowns	Unknowns	Other_	

Description of bird: Size, colors, all field marks OBSERVED. - Unique features of this individual, if any.

A large, very whitish-looking gull. Bill was flesh colored on base with the outer 1/3 rd of the bill dark. Eye was dark, legsspink. Much gray fleckin on the feathers of the back, wing coverts and breast. Wing tips were immaculat whitish. Overall size was at least as large as a Herring Gull, perhaps a little larger. Head appeared proportionately bigger and bill stockier than Herring Gulls nearby. I believe the bird was a 1st winter plumage individual because of the brownish-gray flecking on the body and wing coverts (2cd winter birds are nearly pure white).

Field marks which were <u>not</u> seen. Why not?	Variations from expected field marks.
none	none
Hone .	
	Figure 1
st similar species and describe why or how you eliminated	them.
Iceland is smaller, and typically does winter). Iceland has less stocky look	not have fleshy base to beak (in 1st
Glaucous-winged Gull is browner with a 2cd winter plumage).	ll dark beak in 1st winter plumage (an
zed wineer promogey.	
	The first of the f
op bereit it is to be the control of	. ผู้เด็จ คระไปหน้า และเก็บไป หูแลหางได้เป็น
e e derin al Marinera i incessione	
The second of the second of the second	i di dan Tradicionale interfacioni di la manda de la Propriori di mandale di la companya di dalam Propriori di mandale di la companya di mandale di la companya di mandale di la companya di la companya di la co
**	
abitat at this location. Adjacent or near by habitat, if	significant.
garbage dump	
	net
escribe bird's song or sounds.	Print reporter's name, address, phone number.
	DAVID L. FISCHER
none heard	177 STARCREST
	OREM, UT 84058
•	1 1 0 1 1 1
44	Signature Down & Mount
hat is your experience with this or similar birds?	Corroborating observers not reporting separately.
Have seen many Glaucous Gulls in Mass.	
and 1 in Illinois. Have seen all simi	lar Print Name Signature
species except Glaucous-winged.	
	Print Name Signature
z a	
	Print Name Signature
and the second of the second o	Print Name Signature