

May 15th, 16th, 18th
 Syracuse location
 Probably same bird
 as May 6 & 7

7-1987

- * primaries appear very short beyond tertials
- photo's taken w/ 410 mm lens @ approx. 25 yds.
- underwings not seen
- bird is smaller than the accompanying black bellied
- vocalization not heard

Supercilliaris buffy white

very white

bill appears slimmer & shorter than black plover


Back, crown, & back of neck golden brown mottled w/ black (much less bright than on May 6 & 7th)

face, neck, breast, & belly black

After knowing what to look for & tried to draw the 1st area accurately
 * dark 1st extending only 1st beyond *
 3rd *
 * *

undertail coverts heavily streaked w/ black

Bird is lame (limping) and on 3 observed days stayed w/in 20-50 yds of the road



May 6th 1987

7-1987

7-1987

Impression: bird very close to final breeding plumage

$\frac{3}{4}$ mi. West of
intersection of 3000 W.
& 2700 S. Syracuse
(north side of road)

slightly smaller than accompanying Black Bill
didn't see in flight

Yellowish supercilium
yellow extending down
neck and through
face

Buffy/gray
face mark

Neck ~~turning~~ black (very homogeneous
gray)

whitest just in front of wing
fold

Blackest
on belly


flanks
spotted
or streaked
w/ gray

dark
primaries

Does not represent
true 3rd extension

undertail coverts
streaked w/ black

Back very golden
with black
mottling



May 8th

This is the bird in
adult plumage at this

1/2 mile north of the
intersection of 1800 N. & 4500 W.
location: in Hooper Davis Co.

white supercilium

